

Quark Structure of Light Mesons

Bernard Metsch

*Institut für Theoretische Kernphysik
Universität Bonn
Nußallee 14-16, D53115 Bonn, Germany*

Abstract. On the basis of the Bethe–Salpeter Equation we developed a covariant constituent quark model, with confinement implemented by a linear potential and an instanton induced interaction explaining mass splittings and mixing of pseudoscalar mesons. In addition this interaction yields a scalar (essentially) flavour singlet state at approximately 1 GeV, considerably lower in mass than the corresponding octet states calculated around 1.4 GeV. The validity of the present approach was checked through various electroweak observables. The puzzling properties of scalar mesons is briefly discussed.

A glance at the experimental meson spectrum shows two general features, each with a conspicuous exception: Firstly, states belonging to an orbital angular momentum multiplet exhibit only small spin-orbit splittings: e.g. $f_1(1285)$ and $f_2(1270)$, $a_3(2050)$ and $a_4(2040)$, $K_2(1770)$ and $K_3(1780)$. Exceptional are the low positions of $f_0(980)$ and $a_0(980)$. Secondly, every isovector state has an isoscalar partner almost degenerate in mass: e.g. $\rho(770)$ and $\omega(782)$, $h_1(1170)$ and $b_1(1235)$ up to $a_6(2450)$ and $f_6(2510)$, reflecting the fact that the inter-quark forces are flavor symmetric. An exception is of course the huge splitting of the pseudoscalar mesons $\pi-\eta-\eta'$. Any hadron model should account for this phenomenology, especially in view of the identification of exotics, like hybrids, dimesonic states or glueballs.

The most successful model in this respect is certainly the non-relativistic constituent quark model, where it is assumed that excitations of hadrons are effectively described in terms of constituent quarks interacting through potentials. In a particular version, where confinement was modeled by a linearly rising string-like potential and the widely used Fermi-Breit interaction based on One-Gluon-Exchange was substituted by an instanton induced force [1] one can indeed arrive at a satisfactory description of both meson and baryon spectra. Here, the bulk of states is determined by the confinement potential alone (thus avoiding large spin-orbit splittings) and the instanton induced force selectively acts on pseudoscalar states and accounts for mixing and splitting of the isoscalars. However, this approach can be criticized because binding energies compared to the constituent quark masses can be very large and the Schrödinger wave functions are incorrect at large energies or momentum transfers. Moreover, although the dilepton widths of vector mesons can be accounted for in the non-relativistic approach, see Table 1 weak decay constants are too large by an order of magnitude and the $\gamma\gamma$ -decay results

TABLE 1. Electro-weak meson decays.

Decay	Exp. [2]	RQM	NRQM	Decay	Exp. [2]	RQM	NRQM
$f_\pi[\text{MeV}]$	131.7 ± 0.2	130	1440	$f_K[\text{MeV}]$	160.6 ± 1.4	180	730
$\Gamma_{\pi^0 \rightarrow \gamma\gamma}[\text{eV}]$	7.8 ± 0.5	7.6	30000	$\Gamma_{\rho \rightarrow e^+e^-}[\text{keV}]$	6.8 ± 0.3	6.8	8.95
$\Gamma_{\eta \rightarrow \gamma\gamma}[\text{eV}]$	460 ± 5	440	18500	$\Gamma_{\omega \rightarrow e^+e^-}[\text{keV}]$	0.60 ± 0.02	0.73	0.96
$\Gamma_{\eta' \rightarrow \gamma\gamma}[\text{eV}]$	4510 ± 260	2900	750	$\Gamma_{\phi \rightarrow e^+e^-}[\text{keV}]$	1.37 ± 0.05	1.24	2.06

are even beyond discussion. However, a drastically improved description is found in the relativistically covariant quark model we will now briefly discuss.

The amplitude $\chi_P(x) = \langle 0 | T[\Psi^1(\frac{1}{2}x)\bar{\Psi}^2(-\frac{1}{2}x)] | P \rangle$ is determined by the Bethe-Salpeter equation, which in momentum space reads [3,4]:

$$\chi_P(p) = S_1^F(p_1) \int \frac{d^4 p'}{(2\pi)^4} [-iK(P, p, p')\chi_P(p')] S_2^F(p_2). \quad (1)$$

Here $p_{1/2} = \frac{1}{2}P + / - p$ denote the momenta of quark and antiquark, P is the four momentum of the bound state, S^F is the Feynman quark propagator and K the irreducible quark interaction kernel. Staying as close as possible to the non relativistic potential model we make the following *Ansatz*: The propagators are assumed to be of the free type, i.e. $S_i^F(p) = i/(\not{p} - m_i + i\varepsilon)$, with an effective constituent quark mass m_i ; The kernel K is assumed to depend only on the components of p and p' perpendicular to P , i.e. $K(P, p, p') = V(p_\perp, p'_\perp)$ with $p_\perp := p - (pP/P^2)P$. Integrating in the bound state rest frame over the time component p^0 and introducing the Salpeter (or equal-time) amplitude $\Phi(p) = \int dp^0 \chi_P(p^0, \vec{p})|_{P=(M, \vec{0})}$ we obtain the Salpeter equation:

$$\begin{aligned} \Phi(\vec{p}) = & \int \frac{d^3 p}{(2\pi)^3} \frac{\Lambda_1^-(\vec{p})\gamma^0[V(\vec{p}, \vec{p}')\Phi(\vec{p}')] \gamma^0 \Lambda_2^+(-\vec{p})}{M + m_1 + m_2} \\ & - \int \frac{d^3 p}{(2\pi)^3} \frac{\Lambda_1^+(\vec{p})\gamma^0[V(\vec{p}, \vec{p}')\Phi(\vec{p}')] \gamma^0 \Lambda_2^-(-\vec{p})}{M - m_1 - m_2}, \end{aligned} \quad (2)$$

with the projectors $\Lambda_i^\pm(\vec{p}) = (\omega_i(\vec{p}) \pm H_i(\vec{p}))/2\omega_i(\vec{p})$, the Dirac Hamiltonian $H_i(\vec{p}) = \gamma^0(\vec{\gamma} \cdot \vec{p} + m_i)$ and where $\omega_i(\vec{p}) = \sqrt{m_i^2 + \vec{p}^2}$.

The amplitudes Φ are calculated by solving the Salpeter equation (2) for a kernel containing a confining interaction with a spin structure minimizing spin-orbit effects:

$$\int d^3 p' [V(\vec{p}, \vec{p}')\Phi(\vec{p}')] = - \int d^3 p' v(\vec{p} - \vec{p}') \frac{1}{2} [\gamma^0 \Phi(\vec{p}') \gamma^0 + \Phi(\vec{p}')] . \quad (3)$$

where v in coordinate space is a linearly rising potential $v(|\vec{x}_1 - \vec{x}_2|) = a + b|\vec{x}_1 - \vec{x}_2|$, and 't Hoofts instanton induced interaction acting exclusively on (pseudo)scalars:

$$\int d^3 p' [W(\vec{p}, \vec{p}')\Phi(\vec{p}')] = 4G^{(g, g')} \int d^3 p' w_\lambda(\vec{p} - \vec{p}') [\gamma^5 \text{tr}(\Phi(\vec{p}')\gamma^5) + \text{tr}(\Phi(\vec{p}'))] . \quad (4)$$

Here $G^{(g, g')}$ is a flavor matrix containing effective coupling constants g, g' and w_λ is a regularizing Gaussian, see [4] for details. In order to calculate current matrix elements

in the Mandelstam-formalism [5] we first construct the meson-quark-antiquark vertex function $\Gamma_P(p) = [S_1^F(p_1)]^{-1} \chi_P(p) S_2^F(p_2)$ in the rest frame from the Salpeter amplitude by

$$\Gamma(\vec{p}) := \Gamma_P(p_\perp)|_{P=(M,\vec{0})} = -i \int \frac{d^4 p'}{(2\pi)^4} [V(\vec{p}, \vec{p}') \Phi(\vec{p}')] \quad (5)$$

and then calculate the BS-amplitude for any on-shell momentum P by a boost Λ_P : $\chi_P(p) = S_{\Lambda_P} \chi_{(M,\vec{0})}(\Lambda_P^{-1} p) S_{\Lambda_P}^{-1}$.

The improvement of the results in Table 1 is largely due to inclusion of the second term of equation (2), which is neglected in the non relativistic approach. For other observables we refer to [4,10] concerning spectra, [5,6] for results on form factors, and [7] for an extensive discussion of $\gamma\gamma$ -decays. Here we will only briefly discuss the structure of (pseudo)scalar mesons [8–10]. A comparison of their mass spectra with experimental data is presented in Fig.1. It shows, that the instanton induced interaction not only correctly describes the splitting (and mixing) of the pseudoscalar ground state nonet, but also leads to a particular structure for the scalar mesons: It produces an almost pure flavor singlet state f_0^1 at roughly 1 GeV whereas the flavor octet states f_0^8 , a_0 , K_0^* are almost degenerate at ≈ 1.4 GeV.

This then leads to the following interpretation of experimental data as the scalar $q\bar{q}$ -flavor nonet [8]: the $f_0(1500)$ is not a glueball but the scalar (mainly)-octet meson. The mainly-singlet state could correspond to the broad $f_0(1000)$ -state ($f_0(400-1200)$ of [2]), but there are arguments that it is to be identified with the $f_0(980)$. The isovector and




FIGURE 1. Comparison of the experimental (left side of each column, data from [2]) and calculated (right side of each column) spectrum for (pseudo)scalar mesons. The experimental uncertainty of the resonance position is indicated by a rectangular box, dashed boxes indicate the width.

isodoublet states then correspond to $a_0(1450)$ and $K_0^*(1430)$, respectively. The present model suggests that one isoscalar and one isovector scalar state at 1 GeV is not of the quarkonium type. Indeed, several calculations suggest that these resonances are related to $K\bar{K}$ -dynamics. In this spirit, the $f_0(1370)$ resonance is interpreted as the high energy part of the broad $f_0(1000)$. Furthermore, we could identify the $f_J(1710)$ with the first radially excited scalar state, provided its spin is indeed 0^+ . However, in particular the $f_0(1500)$ was argued to have properties incompatible with a pure $q\bar{q}$ configuration and was suggested to possess a large glue component, mainly because of the suppression of the $K\bar{K}$ decay mode, see [11]. In [9,10] it is shown, that instanton effects can lead to a selective violation of the OZI-rule for decays of scalars into pseudoscalars and that the observed branching ratios can be explained in the framework of a constituent quark model.

We conclude by citing some results on $\gamma\gamma$ decays [7] of scalar and tensor mesons, which constitute a sensitive test of the present approach, and thanking Eberhardt Klempt, Claus Münz, Herbert Petry, Jörg Resag and Christian Ritter.

TABLE 2. $\Gamma(M \rightarrow \gamma\gamma)[eV]$ of scalar and tensor mesons.

M	Exp. [2]	Calc.	M	Exp. [2]	Calc.	M	Exp. [2]	Calc.
$a_2(1320)$	1040 ± 90	734	$f_2(1270)$	2440 ± 300	2040	$f_2'(1525)$	105 ± 17	121
$a_0(1450)$		1390	$f_0(980)$	560 ± 110	1750	$f_0'(1500)$	< 170	161
			$f_0(1370)$	5400 ± 2300				

REFERENCES

1. Blask, W.H., Bohn, U., Huber, M.G., Metsch, B.C., Petry, H.R., *Z. Phys.* **A337**, 327 (1990).
2. Barnett, R.M. *et al.*, (Particle Data Group), *Phys. Rev.* **D54**, 1 (1996) and 1997 off-year partial update for the 1998 edition available on the PDG WWW pages (URL: <http://pdg.lbl.gov/>).
3. Resag, J., Münz, C.R., Metsch, B.C., Petry, H.R., *Nucl. Phys.* **A578**, 397 (1994).
4. Münz, Claus R., Resag, Jörg, Metsch, Bernard C., Petry, Herbert R., *Nucl. Phys.* **A578**, 327 (1994).
5. Münz, Claus R., Resag, Jörg, Metsch, Bernard C., Petry, Herbert R., *Phys. Rev.* **C52**, 2110 (1995).
6. Giersche, Wolfgang I., Münz, Claus R., *Phys. Rev.* **C53**, 2554 (1996).
7. Münz, C. R., *Nucl. Phys.* **A 609**, 364 (1996).
8. Klempt, E., Metsch, B.C., Münz, C.R., Petry, H.R., *Phys. Lett.* **B361**, 160 (1995).
9. Ritter, C., Metsch, B.C., Münz, C.R., Petry, H.R., *Phys. Lett.* **B380**, 431 (1996).
10. Metsch, B.C., Petry, H.R., *Acta Physica Polonica*, **B27**, 3307 (1996).
11. Amsler, C., Close, F.,E., *Phys. Lett.* **B353** 385 (1995); *Phys. Rev.*, **D53** 295 (1996).